GENERAL INDEX TO VOLUME XL

New scientific names of plants and the final members of new combinations are printed in **bold-face** type; synonyms and page numbers having reference to figures and plates, in *italics*; and all other matter in ordinary type.

A

Acanthographis, 311
Acanthotheciopsis, 312
Acasporaceae, 274
Actinoglyphis, 335; *Leprieurii*, 335
Africa, tropical, Some lichens of, 271
Aglaopisma, 389
Agriculture, milpa system of, in Yucatan, 51
Alectoria, 378, 379
Allarthonia, 303
Allarthothelium, 303
Almanac, Bluegrass pasture, 1
Alyxoria, 308
Amphiloma, 389; debanense, 390
Amphiloma, 389
Anaptychia, 399; **Adamesii**, 400; ciliaris, 399; leucomelaena, 399
Anderson, Edgar: The analysis of suspected hybrids as illustrated by Berberis × gladwynensis, 73; and William L. Brown: The popcorns of Turkey, 33
Andrews, Henry N., and Sergius H. Mamay: Some American Medullosas, 183
Animals: dispersal of Persimmon by, 217; observed at Wildwood, 245; relation between bluegrass pastures and, 1, *24*, *26*, *28*
Anomorpha, 312; *turbulenta*, 312
Anthracothecium, 273, 285; cinerosum, 285, 286; Doleschalii, 283; sect. Euanthracothecium, 285; euthelium, 285; guineense, 285, 286; lugescens, 285; sect. Porinastrum, 285; punctuliforme, 285; vitellinum, 285
Antrocarpum, 341
Anzia, 372
Archaeological: and ethnological races of maize, 79, 84; studies of the Maya civilization, 51
Arizona and New Mexico, A contribution to the lichen flora of, 63
Arthonia, 301, 303; applanata, 304; calospora, 305; carneoalbens, 304; cinnabarina, 304, v. dendritica, 304, v. elegantula, 304, v. medusaeformis, 304, v. orbicella, 304, v. reducta, 304, v. rimata, 304, v. speciosa, 304; **elevata**, 305; erythrocarpa, 304, v. roseopallens, 304; fuscescens, 308; gregaria v. dendritica, 304; ilicinodes, 304; leptogramma, 305, 306; **leptogrammodes**, 305, 306; leptographoidea, 304, 305; loangana, 304; microcarpella, 305; **modesta**, 304, 305; palmensis, 304; pertusariella, 304; somaliensis, 304; **subcaesia**, 304, 307
Arthoniaceae, 301, 303, 338
Arthoniopsis, 303
Arthopyrenia, 273, 274
Arthopyrenia sect. *Melanotheca*, 292
Arthopyreniella, 273
Arthothelium, 301, 302, 303, 326; dictyophorum, 302; laceratum, 302
Ascidium, 342; *Cinchonarum*, 342; *rhodostoma*, 343
Asiatic races of maize, cob morphology of, 87, in relation to ethnology and archaeology, 104
Asterisca, 335; *labyrinthica*, 335
Astrotheliaceae, 297
Astrothelium, 297
Aulacographa, 312
Aulaxina, 307

B

Bacidia, 352, 357; abessinica, 358; amylothelia, 358; araneosa, 358; atlantica, 358; byssothallina, 358; decussata, 358; endoleucella v. colorata, 358; sect. Eubacidia, 357, 358; exiguella, 358; farinulenta, 358; golungensis, 358, 359, v. pauciseptata, 359, 360, v. pluriseptata, 359, 360; griseoalba, 358; heteroloma, 360, v. bacteriospora, 358, 361, v. elongata, 358, 361; imitans, 358; infuscata, 358; Kotschyi, 358; leucotrypta, 358; **nigeriensis**, 358, 359; nigrocincta, 358; subg. Scoliciosporum, 274; Scottii, 358; socotrana, 358; stellaris, 358; sublecanorina, 358; submillegrana, 358; trichosperma, 358; vagula, 358; sect. Weitenwebera, 357, 358
Barberry: hybridization of, 73; relationships of two species and their putative hybrids, *74*, *76*, *77*
Bathelium, 275, 278, 280, 286, 288, 296; **compositum**, 289; duplex f. **simplicius**, 289, 290; mastoideum, 289; **papillosum**, 288, 289, 290; **porosum**, 290; **subalbens**, 289, 290
Bathelium, 296
Belonia, 273

Berberis × gladwynensis, An analysis of suspected hybrids as illustrated by, 73
Berberis, 73; × **gladwynensis**, 74, 77, 78; Julianae, 74, 77; verruculosa, 73, 77
Biatora rubina, 355
Birds: list of, eating persimmon fruit, 218; found at Wildwood, 248, 256
Blastenia, 384, 387; albidofusca, 388; albidopallens, 388; Brebissonii v. microspora, 387; cormorensis, 388; lactescens, 388; laetebunda, 388; mauretaniae, 387; maurula, 387; pertenuescens, 387; polioterodes, 388; sinapisperma, 387; sordida, 388
Blasteniaceae, 384
Blastodesmia, 273
Blenothallia, 351
Bluegrass: pasture almanac, 1; pastures, *24-30*
Bombyliospora, 384, 385; cerinella, 385, 386; dominguensis v. intermedia, 385; endoleucites, 385; Meyeri, 385; **nigeriensis**, 385; pruinata, 385; thomensis, 385, 386; **Thoroldi**, 385, 386; togoensis, 385
Borrera, 399
Bottaria, 286, 290; sect. *Anthracothecium*, 285
Brass, L. J., lichens collected by, 271
Brassia, 342
Brigantiaea, 361; *argentea*, 361; *Mariae*, 361; *tricolor*, 361; *tristis*, 361
Brookhill Farm, observations on bluegrass pastures made at, 1
Brown, William L., Edgar Anderson and: The popcorns of Turkey, 73
Bryophagus, 348; *Gloeocapsa*, 348
Buellia, 355, 390; **Adamesii**, 392; afra, 391; albinea, 392; americana v. palmensis, 392; approximans, 392; argilliseda, 392; Caldesiana v. subpruinosa, 391; canescens, 390; Chaudieauiana, 391; cinereocincta, 392; crassa, 391; **Deightoni**, 392, 394; delaevata, 392; sect. Diploicia, 390; sect. Diplotomma, 391; disciformis v. contorta, 391, v. oblongata, 391, 392, v. pachyspora, 391, v. rhodina, 391; dispersa v. cinerascens, 392; sect. Eubuellia, 391; innata, 391; italica v. debanensis, 392; leucina, 391; sect. Melanaspicilia, 391; olivacea, 391; pachyspora, 391; parasema, 390, v. subaeruginosa, 392; paupercula, 392; prosperens, 392; spuria v. ferruginea, 391; stellulata f. subtilis, 392; subalbula, 392; subdives v. mozambica, 392; subimmersa, 391; subpulchella, 391; substigmatea, 392, f. obfuscata, 391; Tobleri, 391; toninioides, 392
Buelliaceae, 390
Bunodea, 280

C

Callopisma, 388
Caloplaca, 384, 388; **arizonica**, 67, *68*; asmarensis, 388; askremensis 389; Balfouri, 390; Beccarii, 390; benguelensis, 388; cinnabarina, 69; citrinella, 389; concinnata, 388; *conglobata*, 389; crispicans, 390; delicata, 390; deplanata, 390; elegantissima, 390; exasperata, 388; flava, 388; flavorubens, 390; granulifera, 390; granuliformis, 390; inconcinna, 388; murorum, 389, 390, v. granuliformis, 390; ochraceofulva, 390; Odoardi, 388; zambesica, 388
Caloplaca, 388
Cameroons, lichens from, 271
Candelaria, 372
Candelariella, 365
Carlosia, 300
Catillaria, 352; subternella, 358
Catinaria, 338, 339, 393
Chapsa, 342; *indicum*, 343
Chicory in pastures, 13, *28*
Chiodecton, 301, 333, 337; **album**, 338; amyloplacoides, 337; amyloplacum, 337; Brunnthaleri, 338; subg. Byssophorum, 301, 337, 338; circumscissum, 337; subg. Enterographa, 337; subg. Euchiodecton, 337; **hypochryseum**, 338; intercedens, 338; irregulare, 337; laceratum, 337; minutulum, 337; molle, 338; mozambicum, 337; nanum, 337; palmensis, 337; rotundatum, 337; sanguineum, 337, 338; seriale, 337; socotranum, 337; sphaerale, 337
Chiodectonaceae, 333
Chiographa, 318
Chlorocyphella, 363
Chromosome knobs on Turkish popcorns, numbers and positions of, 44
Chromosomes: of Delphinium, 132, 134, 164, morphology of, 136, 147; of Diospyros, 213; of Petunia, treated with x-rays, 261
Chroodiscus, 340
Cladoxylon, 198
Clathroporina, 273, 275, 285, 296
Coal balls, fossils of Medullosa found in, 183
Cob morphology among certain archaeological and ethnological races of maize, 79
Coccotrema, 273, 287
Coenogoniaceae, 349
Coenogonium, 349; **congensis**, 350; **Deightoni**, 349; Linkii, 349

Collema, 351; aggregatum, 351; ascaridiosporum, 351; sect. Blenothallia, 351; cheila, 351; sect. Collemodiopsis, 351; conglomeratum, 351; sect. Dicollema, 351; fasciculare, 351; flaccidum, 351; haemaleia, 351; Lactuca, 351; nigrescens, 351, var. minutum, 351; rupestre, 351; *scotinum*, 351; *turgidum*, 351
Collemaceae, 351
Coniochila, 341
Contribution to the lichen flora of Arizona and New Mexico, 63
Corn, see Maize
Coscinedia, 342
Creographa, 327; *brasiliensis*, 327
Crocynia, 301, 397; haematina, 397; Leopoldi, 397
Cryptolechia, 348
Cryptothecia, 301, 302; caesioalba, 302; geniflexum, 302; nigeriensis, 302; Stirtoni, 302; subnidulans, 302; Thoroldi, 302, 303
Cryptotheciaceae, 301, 338
Cyclocarpineae, 338
Cypheliaceae, 300
Cyphelium, 300
Cyphella, 363
Cytological: data of plants showing effect of x-rays, 261; methods used in study of Delphinium, 134; study of persimmon, 213
Cytology, morphology and systematic relationships of Delphinium × Belladonna Hort. ex Bergm., 113

D

Dactylina, 379
Darrow, Robert A., Lichen collection of, 63
Deighton, F. C., lichens collected by, 271
Delphinium, 113; character associations in establishing relationships in species, 125, *127*, *129*, 131; chromosomes of species, 132, 164, of hybrids determined by Propach, 113, 165; cytological study of, 132, *178-182*, of natural species, 136, of derived hexaploids, 150; history of, 113, 132; hybrids of the Belladonna type, 113, 121, *172*, *174*, leaf types in, *113*, pollen studies of, 124, 130; morphology of natural species, 113; satellite expression in species and hybrids, 138; seed germinability in, 155; taxonomy of natural species, 114
Delphinium × Belladonna Hort. ex Bergm., The cytology, morphology, and systematic relationships of, 113
Delphinium, 113; ajacis, 114; azureum, 117, 132; Barlowii, 120, 130; Belladonna, 113, 115, 119, 125; Belladonna type, hybrids of, 117, *118*, 146; × Belladonna Hort., origin of, 113, 117, 119, variability in, 126; Bellamosum, 140; "Blue Butterfly", 121, 124, 130; "Capri", 119; cardinale, 117; cardiopetalum, 132; chamissonis, 116; cheilanthum, 113, 115, 116, *118*, 121, 124, 126, 130, 136, *166*, *170*, var. Chauvieri, 121, var. formosum, 113, var. Hendersonii, 120, var. Middendorffii, 116; ser. Cheilanthoidea, 115; chinense, 121; "Cliveden Beauty", 124, 140, *168*; sect. Consolida, 115; subg. Consolida, 115; consolida, 114; sect. Delphinastrum, 114; sect. Delphinellum, 114; subg. Diedropetala, 115; sect. Elatopsis, 115; elatum, 114, 115, 117, *118*, 119, 121, 126, 129, 130, 132, *172*, × grandiflorum, 121, 133, "Summer Skies", 124, *176*; elatum-splendens, 121, 124; subg. Eudelphinium, 115; formosum, 115, 117, 118, 130, *168*; "Grandiflora," 117; grandiflorum, 114, 115, 117, 119, 123, 126, 130, var. chinense, *118*, 121, 133, 136, *170*, "Blue Butterfly", 121, 122, 137; gypsophilum, 133; Hanseni, 133; Hendersoni, 115, 120, 125, *166*, "Lamartine", *118*, 124, 125, 130; sect. Kolobopetala, 115; Middendorffii, 116, var. parviflora, 116; "Moerheimi", 119, 124, 125, 145; "Mrs. G. Gibson", 117; nudicaule, 117, 133; peregrinum, 114; ser. Racemosum, 115; Ruysii, 133; Smith's Belladonna, *118*, 124, 126, 140, *168*; Staphisagria, 114, 132; sect. Staphisagria, 114; "Summer Skies", 124, *176*, × grandiflorum, 123; tatsienense, 115, 117, *118*; "Tom Thumb", 124; variegatum, 133
Desmaziera, 379
Digraphis, 312; *turbulenta*, 312
Dimerella, 346
Diorygma, 323; *tinctorum*, 323
Diosypros, 211; chromosomes of, 213; history of, 211; discolor, 214; kaki, 214; lotus, 214; texana, 214; virginiana, 211, 214
Diospyros virginiana L., The distribution of, 211; clonal types, 218, *224*, *225*; dispersal by animals and birds, 217, 218; forest associations, 220; in mixed oak associations, *225*; local distribution, 214, *215*, factors affecting, *212*, 214; range of, 211, *212*
Diplogramma, 307, 333
Diplographis, 312
Diploschistes, 312

Distribution of Diospyros virginiana L., 211
Dithelopsis, 273
Dodge, Carroll William: Some lichens of tropical Africa, 271
Dyplolabia, 312; *Afzelii*, 314

E

Ecology: Bluegrass pasture almanac, 1; Distribution of Diospyros virginiana L., 211; Wildwood, a study in historical, 227
Ectographa, 326
Ectolechia, 342, 343
Elmore, Mr. and Mrs. Francis, lichen collections of, 63
Emerson, R. A.: A preliminary survey of the milpa system of maize culture as practiced by the Maya Indians of the northern part of the Yucatan Peninsula, 51
Encephalographa, 393; cerebrinella, 393
Endocoena, 379
Enterodictyon, 333
Enterostigma, 333
Ethnological races of maize, variation in cob morphology among certain, 79
Etter, Alfred G.: Bluegrass pasture almanac, 1; Wildwood—a study in historical ecology, 227
Euthelenella, sect. *Microglaena*, 275
Evernia, 379
Everniopsis, 379

F

Farriola, 300
Fissurina, 312; *Dumastii*, 312; *incrustans*, 312
Florissant, Mo., history of an estate near, 227
Flower abnormalities of plants treated with x-rays, 260
Forests in the St. Louis area, 1817-18, 230
Fossils: of Medullosa, 183; of persimmon, 211
Fouragea, 307
Fulgensia, 385
Fungi of tropical Africa, 401

G

Gage, Marilyn Amy: The cytology, morphology and systematic relationships of Delphinium × Belladonna Hort. ex Bergm., 113
Gasparrinia, 385, 389; murorum, 389
Glaucinaria, 323
Glyphis, 333, 334; cicatricosa, 334, var. pulvinata, 334; *labyrinthica*, 334, 335; *tricosa*, 334
Gophers in bluegrass pastures, 3, *30*
Graphidaceae, 311, 333
Graphidula, 318
Graphina, 311, 312, 318, 323; abstracta, 324; aethiopica, 324; ambrizensis, 324; arthothelioides, 324, 325; Brunnthaleri, 324; carneum, 323; *chlorocarpa*, 312; confertum, 323; Deightoni, 324; heterospora, 324; *incrustans*, 312; javanica, 318; Pelletieri v. macrior, 324; sect. Platygrammina, 326; sect. Platygrammodes, 311, 325; sect. Platygraphina, 326; plicatum, 323; Poitiaei, 327; pyrenuloides, 324; raddacensis, 323; rudescens, 324; serpentarium, 323; socotrana, 324; sorediella, 324, 328; straminea, 324; subhiascens, 324; turgidum, 323; ulcerata, 324, 325; varians, 324
Graphis, 308, 311; Afzelii, 312, 314; albonotata, 312; aperiens v. pruinosa, 313; aterrima, 314; betuligna, 312; brachycarpa, 313; caesia, 312; *caribaea*, 303, 318; Cerasi, 312; chlorocarpa, 312; *chrysenteron*, 327; *cicatricosa*, 334; *coccinea*, 318; comma, 313; *deducta*, 330; Deightoni, 314, 316; *dendritica*, 318; Dracaenae, 313; elegans, 312; erythrocardia, 314; exalbata, 313; guineensis, 314, 317; hololeuca, 323; hyalinella, 313; increbrior, 313; infida, 313; inusta, 323; Junghuhnii, 323; labyrinthica, 335; Limae, 313; *lynceodes*, 322; mocimbensis, 313; *Montagnei*, 327; myriocarpoides, 313; myriocarpiza, 313; nigeriensis, 313, 315; nivea, 314; ochroplaca, 313; ondensis, 313, 316, 318; oxyclada, 313, 314; pallescens, 313; palmensis, 313; Poitiaei, 323; polymorpha, 327; pulverulenta, 312; raddacensis, 323; roseotincta, 312; rufula, 312; *scalpturata*, 318, 327; Schroederi, 313; subg. *Scolaeospora* sect. *Glyphis*, 334; scripta, 311, 312; serpentina, 312; striatula, 313, var. cuspidata, 313; subintricans, 313; superans, 313; tenella, 314; Thoroldi, 313, 336; timidula, 312; triticella, 313; turbulenta, 312; *ulcerata*, 325
Grazing habits of stock, effect of, on grasses, 2
Gyalecta, 346; *plurilocularis*, 348; sect. *Secoliga*, 348
Gyalectaceae, 346
Gymnographa, 307
Gyrostomum, 340, 346

H

Haematomma, 331, 365, 370; Brassii, 370; dactylopholis, 370, 371; similis, 370; subarthonioideum, 370; vulgare, 370
Hagenia picta v. rupicola, 397

Harlan, Dr. Jack, collection of Turkish popcorns made by, 33
Harpidium, 365
Helminthocarpon, 307; pruinosum, 318
Hemithecium, 323; *chrysenteron*, 323
Henry, Mrs. J. Norman, hybrid barberry in garden of, 73
Heterangium, 198
Heterodermia, 399
Heterothecium, 361; *Berteroanum*, 361
Holothelis, 273
Horse grazing, effect of, on bluegrass pastures, 7
Huea, 384, 387
Hybridization in the evolution of natural populations, 73
Hybrids: An analysis of suspected, as illustrated by Berberis × gladwynensis, 73; of Delphinium Belladonna Hort. ex Bergm., 113; of Petunia, treated with x-rays, 259
Hymenodecton, 318
Hypochnus, 337; *rubrocinctus*, 337
Hysterina, 308

I

Icmadophila, 365
Imbricaria, 372
Ionaspis, 346, 347; **ascidioides**, 347
Irradiation, effect of, on Petunia, 259

K

Kansas: coal-ball fossils found in, 184, 189
Kuttlingeria, 385

L

Larkspurs, Garland, see Delphinium
Lathagrium, 351
Laurera, 275, 280, 286, 296; astroidella, 296; elegans, 296; madreporiformis, 296; **marginata**, 296; **nigeriensis**, 296, 297; ochroleucodes, 296; pauperrima, 296; velata, 296
Leaf abnormalities of plants treated with x-rays, 260, morphology of, 265, *268*, *270*
Lecanactidaceae, 338
Lecanactis, 331, 338, 339; amylacea, 339; *confluens*, 318; **Deightoni**, 339; flava, 339; flavescens, 339; *leucophora*, 339; *lobata*, 339; *Montagnei* subsp. *deducta*, 340, subsp. *deducens*, 340
Lecania, 365; sect. *Platylecania*, 385
Lecaniopsis, 346
Lecanora, 355, 365; aequinoctialis, 367, 368; africana, 367, *367*; subg. Aspicilia, 348; **Brassii**, 366; callimorpha, 367; carneolutea, 348; carneosulphurea, 366; *conglobata*, 389; corallina, 367; *dactylopholis*, 371; ecoronata, 367; elatina, 370; **elmorei**, 65, *66*; subg. Eulecanorea, 366; fibrosa, 367; flavidonigrans, 367; flavidocarnea, 366; flavochracea, 367; *granifera*, 354; granulescens, 367; *haematomma*, 370; homaloplaca, 367; latericola, 367; laterigena, 366; lateritica, 366; leprosa v. carneola, 366, v. rufocarnea, 366; leucoplaca, 367; Monodorae, 366; notha, 367; sect. *Ochrolechia*, 369; pedata, 367; pleospora, 366; poliothallina, 366; polytropoides, 366; rubiniza, 367; sabulosa, 367; Sanctae-Helenae, 366; socotrana, 367; subcongruens, 367; subfusca v. ferax, 367, v. glebulosa, 367, v. leucoblephara, 367, v. melaleuca, 367; triguttulata, 367; tropica, 367; usambarensis, 366
Lecanoraceae, 365
Lecidea, 352, 353; albocincta, 354; amaniensis, 354; angolensis, 354, v. orientalis, 354; argillicola, 354; subg. Biatora, 353, 355; bogosensis, 354; *Buettneri*, 364; Cagnii, 354; caliginosa v. rhacocarpa, 354; capopensis, 354; *carnea*, 357; carneorufa, 354; cinereorubra, 353; citima, 354; endochrysea, 354; subg. Eulecidea, 353; furfurosula, 353; glauconigra, 354; *golungensis*, 359; granifera, 353, 354, v. lecanoroides, 355, v. leucotropoides, 355; gyalectoides, 354; *heteroloma*, 360; hypomeloides, 354; leptobola, 354; lithagoga, 354; microspermoides, 353, 356; mossamedana, 354; **nigeriensis**, 354, 356; nigricans, 354; olivacea v. alsodinea, 361; pannosa, 353; phaeophthalma, 354; rosella, 357; rubina, 354, 357; Sangeana, 354; sorediata, 395; tenuis, 353, 355; trachytica, 354; tuberculosa, 385; xanthinula, 354
Lecideaceae, 352, 384
Leightonia, 288; *porosa*, 290
Leiogramma, 318; *pruinosum*, 318; *punctiforme*, 318; *sericeum*, 318
Leiorreuma, 318, 327; *Lyellii*, 318; *tartareum*, 318
Leptogium scotinum, 351; turgidum, 351, 353; vesiculosum, 351
Leptorhapis, 273
Leptotrema, 340
Lethagrium, 351; *ascaridiosporum*, 351; *conglomeratum*, 351; *fasciculare*, 351; *rupestre*, 351; *turgidum*, 351
Letharia, 379
Leucogramma, 323, 326; *carneum*, 323; *confertum*, 323; *plicatum*, 323; *raddacense*, 323; *serpentarium*, 323; *turgidum*, 323

Lichen crispus, 351; floridus, 379; haemostomma, 370; lepadinus, 341; *lynceus*, 339; pallescens, 369; puniceus, 370; ruber, 348; rugosus, 308; saxatilis, 372; scriptus, 308, 312; speciosus, 400; ventosus, 370
Lichen flora of Arizona and New Mexico, A contribution to the, 63
Lichens of tropical Africa, Some, 271
Limboria, 312; *constellata*, 312; *tridens*, 312
Lithographa, 307
Lithothelium, 297
Lopadium, 352, 361; **Deightoni**, 361, 362; glaucophaeoides, 361; lecanorinum, 361; Newtonianum, 361; **nigrobrunneum**, 361, 363; pezizoideum, 361; **sepiaceum**, 361
Loxospora, 370

M

McQuade, Henry A.: Some x-ray effects in Petunia, 259
Maize: Archaeological remains of, in North America, 84, 86, in South America, 87; cob morphology of, 79; milpa system of culture as practiced by the Maya Indians of Northern Yucatan, 31; Popcorns of Turkey, 33; present-day races of, in Asia, 87, in North America, 83, in South America, 87; significance of popcorns in history of, 33; Variation in cob morphology among certain archaeological and ethnological races of, 79
Malvern B. Clopton Experimental Farm at Clarksville, Mo., 1
Mamay, Sergius H., and Henry N. Andrews: Some American Medullosas, 183
Manure, animal, in pastures, 2
Maya Indians, in northern part of Yucatan, Maize culture as practiced by, 51
Mazozia, 323
Medullosa, 183; anglica, 184, 197, var. ioensis, 186, var. thiesseni, 186; subg. anglorota, 186; distelica, 184, 186, *187*, *203*, *205*, *209*; elongata, 186; endocentrica, 186, 188, 197; **grandis**, 191, *192*, *193*, *195*, 197, *200*, *205*, *207*, *208*; heterostelica, 183, 184, *187*, 188, *190*, 191, 196, *205*, *206*; leuckarti, 198; noei, 191, 196, *209*; olseniae, 198; primaeva, 186, 188; solmsi, 198; thompsonii, 184, *185*, 188, 196, *202*, *203*, *204*
Medullosas, Some American, 183
Medusulina, 333
Megalographa, 326; *hysterina*, 327
Megalopsora, 372
Megalospora, 352
Meissneria, 296; *varia*, 296
Melampidium, 338
Melanodecton, 337; *indicum*, 337
Melanotheca, 281, 286, 290, 292; Achariana, 291, 292, 295; *Achariana* var. *angolensis*, 293, var. *obscurascens*, 295; **angolensis**, 293; **cameroonensis**, 293, 295; **nigeriensis**, 282, 293; **obscurascens**, 293; **porosa**, 293, 294; purpurascens, 292; **pusilla**, 292, 296
Melaspilea, 311, 318
Menegazzia, 372
Meristosporum, 296
Micrographa, 307
Microphiale, 346
Milpa system of maize culture, as practiced by the Maya Indians of northern Yucatan, 53
Minksia, 333
Monoblastia, 273, 287
Morphology: of Delphinium × Belladonna Hort. ex Bergm., 113; of Petunia plants treated with x-rays, 265; of the cob, variation in, among certain archaeological and ethnological races of maize, 79
Mycoblastus, 352
Mycoglaena, 274, 275
Myriostigma, 302; *candidum*, 302
Myriotrema, 342; *album*, 342; *olivaceum*, 342
Mytilidium, 401
Myxodictyon, 365
Myxotheca, 302; *hypocreoides*, 302

N

New Mexico, Arizona and, A contribution to the lichen flora of, 63
Nickerson, Norton H.: Variation in cob morphology among certain archaeological and ethnological races of maize, 79
Nigeria, lichens from, 271
North American races of maize, 105; cob morphology of, 83
Nyasaland, lichens from, 271

O

Ocellis, 342
Ocellularia, 340, 342; albescens, 343, 346; sect. *Ascidium*, 342; subg. Ascidium, 347; cavata, 343, 345; discoidea, 343; fumosa, 343; microspora, 344; mozambica, 343; myriospora, 343; sect. *Myriotrema*, 342; obturata, 342, 343; Poncinsiana, 343; porinoides, 343; **scolecospora**, 335, 343, 345; subterebrata, 343; terebrata, 345; **trypanea**, 343, 344
Ochrolechia, 365, 369; **palmicola**, 369
Omastin, H. A., lichens collected by, 271
Opegrapha, 307, 308; alboatra, 309; albocinerea, 309; *astroidea*, 339; aterula, 309; cretacea, 309, 327; cylindrica, 308; delicata, 309; depressum, 327; diophora, 308;

Dracaenarum, 309; elegans, 309; sect. Euopegrapha, 308; *faginea*, 308; frustulosa, 310; graphidiza, 309, 310, 311; humilis, 309; *illecebrosa*, 339; lepidella, 309; leptographa, 309; lichenoides, 308; loandensis, 309; *Lyellii*, 318; medusulina, 309; Menyhartii, 309; microspora, 309; mozambica, 310; **nigeriensis**, 309, 310; notha, 308; sect. Pleurothecium, 307; *Poitiaei*, 327; prosedoides, 309; prosodea, 309, 311; radiata, 303; rufa, 309; septemseptata, 309; sordida, 327; sororiella, 309; *streblocarpa*, 327; subcalcarea, 309; subgraphidiza, 310; subnothella, 309; tartarea, 318; Vainioi, 308; vermelhana, 309; vestita, 309; viridulata, 309
Opegraphaceae, 307, 333, 338
Ophioparma, 370
Oropogon, 378, 379
Oxystoma, 308; cylindrica, 308

P

Pachyphiale, 346
Paleobotany, 183
Pannoparmelia, 372
Parathelium, 282, 300
Parmelia, 372; abessinica, 375, f. glabrior, 375, f. sorediosa, 375; africana, 375; amaniensis, 375; sect. Amphigymnia, 374; asmarana, 373; caperata v. isidiophora, 375; cetrata v. subisidiosa, 373; coilocarpa, 373; concrescens, 373; coronata v. denudata, 373; ducalis, 374; ecaperata, 375; eurycarpa, 375; Hanningtoniana, 373; Hildenbrandtii, 375, v. ciliata, 375; homolotera v. bryophila, 373, v. hypocraea, 374; hypocrysalea, 375; subg. Hypogymnia, 372; sect. Hypotrachyna, 373; incisa, 397; isidiza, 374; laevigatoides, 373; latissima, 378; leptasca, 373; leucorhiza, 374; lobulascens, 375, v. isidiosissima, 374, 375, 377; Mangenoti, 373; Menyhartii, 375; modesta, 375; neirobiensis, 375; *nilgherrensis* v. *subciliaris*, 377; nitens, 375, f. isidiosa, 374; olivetorum v. esorediata, 375, v. sorediosa, 375; **orchidophila**, 373, 374; pedicellata, 375, v. subbullata, 375; sect. Physcioideae, 372; procera, 375; pseudotinctorum, 374, 375; revoluta v. ambigua, 373; Schimperi, 375; Schweinfurthii, 373, 375; Scottii, 373; sensibilis, 373; somaliensis, 375; Soyauxii, 375, 378; subcetrarioides, 375; **subciliaris**, 375, 377; tenuirima f. sorediata, 373; tiliacea v. hypoleuca, 373, v. rimulosa, 373; uberrima, 375; usambarensis, 373; vicinior v. bryophila, 373; zambesica, 375
Parmeliaceae, 372
Parmentaria, 297; Chevalieri, 286
Pastures, Bluegrass, 1, *24*, *26*, *28*, *30*
Persimmon: distribution of the native, 211, by animals and birds, 217, 218, factors affecting, 214; habitat types, 218, *224*
Pertusaria relata, 345
Pertusariaceae, 287
Petunia, Some x-ray effects in, 259; cytological data, 261; morphological abnormalities, 260, *268*, *269*, *270*; pollen counts, 263
Phaeographina, 312, 326; **alata**, 328, 332; Balfourii, 327; brasiliensis, 327; caesiopruinosa, 330, v. striolata, 327; ceramia, 327; chrysenteron, 327; deducta, 327, 330; **Deightoni**, 327, 329; fuscescens, 328; **innata**, 327, 328; **leptotremoides**, 328, 331; **leucophora**, 327, 329; Montagnei, 327; mozambica, 327, 330; paucilocularis, 328; pezizoidea, 327; quassiaecola, 327; scalpturata, 318, 327; scriptitata, 327, 329; streblocarpa, 327; suffultum, 318
Phaeographis, 311, 312, 318; confluens, 318; **Deightoni**, 319, 320; dendritica, 318; dendriticella, 319; duplicans, 319; flammula, 318; inusta, 323, v. radians, 319; leucothallina, 319; lignatilis, 319; Lyellii, 318; lynceodes, 284, 319, 322; medusulina, 318; micrograpta, 319; micrograptoides, 319; navicularis, 319; **ochracea**, 319, 320; palmarum, 319; paragrapta, 319; Phyllocharis, 319; platycarpa, 319; punctiformis, 318; sericea, 318; sexlocularis, 319; **sierraleonensis**, 319, 321; subdevelans, 319; subnivescens, 319; **tecta**, 319, 322; tigrinella, 319; tridens, 312
Phaeotrema, 340
Phialopsis, 348
Phlyctella, 365
Phlyctidia, 365
Phlyctis, 365
Phragmopyxine, 395
Phyllophthalmara, 340
Phyllopsora, 364; brachyspora, 364; Buettneri, 364; pannosa, 364; parvifolia, 364
Phyllopsoraceae, 364
Physcia, 396; abbreviata, 397, 398; aegiliata, 397; afra, 397, 398; africana, 397; areolata, 397; adscensionis, 397; ascensionis, 397; dilatata, 397; endopyxinea, 397; flava, 397; isidiophora, 397; obscurella, 397, v. fusca, 397; palmarum, 397; picta v. coccinea, 397, v. flavicans, 397; Poncinsii, 397; reticulata, 397; setosa f. vitellina, 397; singularis, 397

Physcia, 372
Physciaceae, 395
Physcidia, 372
Pietzschea, 198
Pioneer Hi-Bred Corn Co., Turkish popcorns grown at, 33
Platygramma, 318; *dendriticum*, 318; *Lyellii*, 318; *suffultum*, 318
Pliarona, 326, 327; *Montagnei*, 327
Pollen of Petunia plants, effect of x-rays on, 261, 263
Polyblastia, 274
Polyblastiopsis, 272, 273, 274, 275, 280, 296; **fulva**, 275; haematochroa, 275; **linearis**, 275, 276; **pyriformis**, 275, 276; **sphaerica**, 275, 277; tropica, 275
Polystroma, 340
Popcorns of Turkey, 33
Popcorns: Turkish, 33, *46*, *48*; chromosome knobs of, 44; distribution of, *40*; list of types, 42; variations in, 34, *38*; Aegean race, *37*, 39; Asiatic race, 35, 46, *48*
Porina, 273, 278, 292; compuncta, 292
Porodothion, 292; *arthonioides*, 292
Porothelium, 292, 300; *arthonioides*, 292
Prairies around St. Louis in 1817, 130
Precipitation, effect of, on distribution of Diospyros virginiana, 213
Preliminary survey of the Milpa system of maize culture as practiced by the Maya Indians of the northern part of the Yucatan Peninsula, 51
Protoblastenia, 384
Pseudolecanactis, 338
Pseudophyscia, 399
Pseudopyrenula, 272, 273, 278, 279; bengoana, 278, 280; *composita*, 289; conica, 278; **Deightoni**, 275, 278, 279; diluta, 280; diremta, 280; *duplex* f. *simplicior*, 290; elliptica, 280; flavicans, 280; sect. Hemithecium, 280; sect. Homalothecium, 279; sect. *Holothecium*, 279; infossa, 278, 280; polyphragmia, 278; sect. Pseudacrocordia, 281; subgregaria, 280; superans, 280; subg. *Trypethelium* sect. *Bathelium*, 289, sect. *Eutrypethelium*, 290
Psora Buettneri, 364
Psorella, 365
Psoromidium, 364; *Wellingtonii*, 364
Psorothecium sect. *Bombyliospora*, 385
Ptychographa, 307, 333
Pycnographa, 333
Pyrenastrum, 297; **erumpens**, 298; **parathelioides**, 298, 299; **plicatum**, 297; **pruinosum**, 298; septicolare, 297
Pyrenodesmia, 385, 388; **conglobata**, 388, 389
Pyrenopsis haemaleia, 251
Pyrenula, 273, 278, 280, 342; Acaciae, 281; Alni, 281; annularis, 279; aspistea, 282; ceratina, 279; discolor, 279; discolorella, 279; eucalypta, 281, 283; Euphorbiae, 281; fuscolurida, 281, 283; sect. Fusidospora, 281; gibberosa, 281; glabriuscula, 282; Gravenreuthii, 281, 292; heteroclita, 282, 284, v. *denigrata*, 284, v. *minuscula*, 282; Limae, 282; mamillana, 282, 284; mastophorizans, 281; subg. *Melanotheca*, 292; mozambica, 281; neglecta, 279; nitida, 281; *obscurascens*, 295; obscurata, 281; ocellata, 281; oculifera, 281; parva, 281; porinoides, 279; Pupula, 279; sect. Pseudoacrocordia, 281; subaperta, 281; trombetana, 281, 282; *trypanea*, 344
Pyrenulaceae, 272, 333
Pyrgillus, 300
Pyrochroa, 318; *flammula*, 318; *javanica*, 318; *medusulina*, 318
Pyrrhographa, 318; *flammula*, 318; *javanica*, 318; *medusulina*, 318
Pyxine, 395; **Adamesii**, 395, 396; Chevalieri, 395; chrysantha, 395; Cocoes v. chrysantha, 395, v. congensis, 396, v. convexior, 396, v. sorediigera, 395; convexa, 396; devertens, 395; dimorpha, 395; endoleuca, 395; Meissneri v. sorediosa, 395; retirugella f. isidiigera, 395; sulphurans, 395

R

Racodium, 349
Radiation effects on Petunia: cytological, 261; on flowers, 260, *270*; on leaves, 260, 265, *268*, *270*
Ramalina, 378, 379
Ramonia, 346
Rhizocarpon, 352
Riddlea, 286, 287; **papillosa**, 287
Rinodina, 355, 390; **darrovii**, 69, *70*; *nimbosa*, 71; phaeocarpa, 71
Rudolph, Emanuel D.: A contribution to the lichen flora of Arizona and New Mexico, 63

S

Sagiolechia, 346
St. Louis area: animal population at Wildwood in, 215; bird population in, 248, 256; occurrence of Diospyros in, 214, 215, 216; prairie and timber area in 1817, 230; plant species found in, 254; tree species in, 232; Wildwood, 227

Sarcographa, 333, 335, 393; Cascarillae, 335; Cinchonarum, 335; sect. Eusarcographa, 335; labyrinthica, 284, 335; Thoroldi, 335, 336; tigrina, 335
Scaphis, 308; anfractuosa, 308; compressa, 308
Schismatomma, 338
Schistophoron, 300
Schistostoma, 341; dehiscens, 341
Sclerographis, 307, 401
Sclerophyton, 333
Secoliga, 346, 348; **Deightoni**, 348; **pluriloc**ularis, 348
Segestrella sect. *Melanotheca*, 292
Semigyalecta, 346
Sierra Leone, lichens from, 271
Siphula, 379
Skallerup, Harry R.: The distribution of Diospyros virginiana L., 211
Solenographa, 318
Solenospora, 365
Solorina, 303
South American races of maize, 35, 86
Sphaerophoropsis, 352
Sporopodium, 363
Stegobolus, 342; *Berkeleyanus*, 343
Steloxylon, 198
Stenographa, 323
Stirtonia, 301, 311
Stirtoniopsis, 301, 311
Stock, grazing habits of, 2
Sutcliffia, 197
Swine, effect of excretory behavior on bluegrass, 8
Sychnogonia, 273; *Bayrhoefferi*, 273
Synarthonia, 303
Syncesia, 337; *albida*, 337
Synechoblastus, 351
Systematic relationships of Delphinium × Belladonna Hort. ex Bergm., 113

T

Temperature, effects of, on distribution of Diospyros virginiana, 213
Thalloloma, 323
Thamnolia, 379
Thecaria, 326; *quassiaecola*, 327
Thecographa, 326; *ceramia*, 327
Thelenella fulva, 275; sect. *meristosporum*, 296
Thelocarpon, 273; sect. *Thelopsis*, 273
Thelococcus, 273
Thelographis, 326
Thelopsis, 273; flaveola, 273; inordinata, 273; rubella, 273; **selenospora**, 273, 274; subporinella, 273
Theloschisma, 318; *Eschweileri*, 318
Thelotrema, 340, 341; **cameroonensis**, 341, 342; *cavatum*, 345; foratum, 341; *microporum*, 343; *obturatum*, 342; subg. *Ocellularia*, 342; Pechuelii, 341; *porinoides*, 343; schistostomoides, 341; *subterebrata*, 343; terebrata, 344; sect. Tremotylopsis, 342; turgidulum, 341; *urceolare*, 342; varioloides, 341
Thelotremaceae, 340
Theobroma of tropical Africa, lichens growing on, 271
Thorold, C. A., lichen collection of, 271
Tomasellia, 286
Toninia, 352
Trees at Wildwood, St. Louis Co.: early descriptions, 230; in 1946, 232, 254; ages of, 234, diameters, heights and growth rates, 234, compared to witness trees, 237, density of, 239; vegetation beneath, 245
Tremotylium, 340; africanum, 340; angolense, 340
Trichophyma, 303
Triophthalmidium, 385
Trypetheliaceae, 286, 333
Trypethelium, 286, 290; **aciculare**, 290, 291; anomalum, 290, 291, 292; *anomalum* f. *leucostomum*, 292, v. *obscurascens*, 295; sect. *Bathelium*, 288; coccinatum, 290; *compositum*, 289; conglobatum, 292; *duplex* f. *simplicius*, 290; Eluteriae, 290; sect. *Eutrypethelium*, 290; **leucostomum**, 290, 292; marginatum, 289; *mastoideum*, 289; meristosporum, 290; *papillosum*, 288, 290; Perrotetii, 292; *platystomum* f. *leucostomum*, 292; *porosum*, 289, 290; *pusillum*, 296; pustulatum, 290; sphaerocephalum, 290; *Sprengelii* v. *porosa*, 290; *subalbens*, 290
Turkey, Popcorns of, 33
Turkish popcorns, 33; chromosome knobs, 44; distribution of, 40; grown in Iowa, 33; variations in, 34, 38; varieties of, 42, Aegean race, 27, Asiatic race, 36, *46*, *48*
Tylophorella, 300
Tylophoron, 300; ascidioides, 300; moderatum v. modestius, 300

U

Uganda, lichens from, 271
Urine, animal, effect of, on pasture grasses, 2, 9, *24*, *26*, *28*, *30*
Usnea, 379; abissinica, 382; acanthera, 382; aequatoriana, 381; africana, 383; albomaculata, 382; amplissima, 383; arguta, 383; *barbata* v. *florida* f. *australis*, 383;

Baileyi, 380; bicolorata, 382; Bornemuelleri, 382; cartilaginea, 381; chloreoides, 383; complanata, 380; contorta, 383; corrugata, 380; cristata, 380; decipiens, 381; distensa, 382; elata, 380; erubescens, 382; subg. Eumitria, 380; subg. Euusnea, 380; exasperata, 381; firmula, 380; flaccida, 381; flavescens, 380; Flotowii, 382; subsp. serpentaria, 382; fusca, 381; gigas, 383; gracilis, 381; Haumanii, 381; hispida, 382, 383; hispidula, 381; implicata, 380; incrassata, 381; Ledienii, 382; leprosa, 380, 383; Liechtensteinii, 380; livida, 383; Lyngei, 383; Meyeri, 382; moniliformis, 381; myrioclada, 382; obtusata, 382; ochrophora, 381; perhispidella, 380; perspinosa, 381; picta, 382; praelonga, 380; pseudocyphellata, 380; pulchella, 381; pulverulenta, 380; rugosa, 380; ruvidescens, 381; ruwenzoriana, 381; saxatilis, 381; scutata, 382; simplicissima, 381; speciosa, 380; subflorida, 382; submollis, 381; terrestris, 380; torrida, 382; trachyna, 382; trichodeodes, 383; umbrata, 383; undulata, 381; usambarensis, 381; Vainioana, 381; versicolor, 380; vesiculata, 380; Welwitschiana, 380
Usneaceae, 378
Ustalia, 303; *anguina*, 323

V

Variation: in cob morphology among certain archaeological and ethnological races of maize, 79; in Turkish popcorns, diagrams illustrating, *34*, *38*
Verrucaria *aspistea*, 318; *mamillana*, 284, 289; *guineensis*, 285; *parella*, 369; *porosa*, 290; *trypanea*, 344; *subcaerulescens*, 275
Volvaria, 341

W

Weeds: control of, in maize culture, 58; in bluegrass pastures, 13, 16, *28*
Wildwood—A study in historical ecology, 227: animals inhabiting, 245; birds and bird nests observed at, 248, 249, 256; history of area, 227; prairie and timber areas in 1817, 230; plant species found at, 254; tree population in 1946, 232, ages of, 235, sizes, 234; vegetation beneath trees, 246

X

X-ray: effects in Petunia, Some, 259; methods of irradiation, 259; visible effects, 260
Xanthocarpia, 384
Xenocladia, 198
Xylastra, 308
Xylographa, 307
Xyloschistes, 312

Y

Yucatan Peninsula, Maize culture as practiced by the Maya Indians in the northern part of the, 51

Z

Zea Mays, see Maize
Zwackhia, 308; *involuta*, 308

Vol. 40,

s, dia-

, 284,
porosa,
s, 275

58; in

cology,
ds and
, 256;
timber
found
, 232,
on be-

259;
ble ef-

s prac-
north-